



Lawrence Livermore National Laboratory

## **"Deuterium Hugoniot up to 120 GPa (1.2 Mbar)" (U)**

**UCRL-PROC-201017**

**W. J. Nellis,\* R. F. Trunin,\*\* A. I. Bykov,\*\* and R. I. Il'kaev\*\***

\*Lawrence Livermore National Laboratory, Livermore, California 94550;

\*\*All-Russia Research Institute of Experimental Physics, Sarov, Russia 607190

December 1, 2003

Submitted to the Nuclear Explosives Design Physics Conference  
Los Alamos, New Mexico,  
October 19-24, 2003

## Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Deuterium Hugoniot up to  
100 GPa (1 Mbar) Pressures

W. J. Nellis  
Lawrence Livermore National Laboratory


R. F. Trunin, A. I. Bykov, R. I. Il'kaev  
All-Russia Research Institute of Experimental Physics  
(VNIIEF), Sarov  
(Formerly known as Arzamas-16)

UCRL-POST-200229

Work performed under the auspices of the U.S. Department  
of Energy by Lawrence Livermore National Laboratory under  
Contract W-7405-Eng-48.

This presentation is unclassified.

## The Controversy: Which reported deuterium Hugoniot is correct?




On the left is the Hugoniot measured by Knudson et al at Sandia Z Machine and numerous theories: (Sesame (Kerley), Tight-binding MD (Lee Collins), etc.

On the right is Hugoniot measured by Da Silva et al at LLNL Nova Laser and Ross' prediction.

To resolve which is correct, independent accurate measurements are needed. Arzamas has a 50-year history of doing high-quality Hugoniot measurements.

Arzamas started measuring the deuterium Hugoniot with partial support from LLNL to resolve this issue.

VNIIEF (Arzamas-16) experiments




Steel shell has maximum velocity of  $\sim 15$  km/s.

Corrections are made for hemispherical convergence.

$\sim 5$  shots are done at each pressure to get good statistics.

Experimental data for shock velocity  $u_s$  versus particle velocity  $u_p$  for liquid deuterium




Solid circles: Nellis and Mitchell (LLNL two-stage gun)

Solid squares: Knudson et al (Sandia Z Machine)

Open triangles: Trunin et al (Arzamas HE-driven converging shocks)

Open squares: Da Silva et al (LLNL Nova Laser)

Hugoniots of liquid deuterium



Solid curve: calculated from fit to  $u_s - u_p$  data of Sandia and Arzamas


Dashed curve: calculated from fit to Nova  $u_s - u_p$  data

Error bars at 100 GPa: uncertainties in fits due to uncertainties in measured  $u_s$

Dotted curve: PIMC calculation of Militzer and Ceperley, including indicated temperatures

Open circles: *ab initio* calculations of Desjarlais (Sandia)

Slopes of  $u_s(u_p)$  versus  $u_p$  for liquid deuterium



Solid curve: calculated from fit to  $u_s - u_p$  data of Sandia and Arzamas

Dotted line: calculated from PIMC results

Open circles: calculated from *ab initio* results

Dot-dash: values for monatomic and diatomic ideal gases and free electrons, as indicated

Dashed line: calculated from fit to Nova  $u_s - u_p$  data

Conclusions

1. Deuterium Hugoniot data measured by Sandia and Sarov are in excellent agreement from 30 to



100 GPa (0.3 to 1 Mbar). The error bars on these data are  $\sim 1/3$  those of the LLNL Nova Laser.

2. This controversy is resolved by the meaning of term “error bar”. The “true” Hugoniot lies within the Nova error bars. The Sandia/ Sarov data tells us the “true” Hugoniot lies at the “stiff” end of Nova error bars.
3. Signature of molecular dissociation between 20 and 50 GPa is a relatively shallow ( $\sim 10\%$ ) minimum in the slope  $S = du_s/du_p$ .
4. Slope  $S = du_s/du_p$  cannot resolve whether or not there are any interactions between particles at Mbar shock pressures ( $\sim 1$  eV); i.e.,  $S$  does not resolve the difference between PIMC with interactions and free electrons without interactions.
5. Kinetic energy dominates potential energy.

Bottom line

The Hugoniot of Sandia/Sarov is correct.